



PRIMARY RESEARCH ARTICLE

Global Change Biology WILEY

Uncovering hidden genetic variation in photosynthesis of field-grown maize under ozone pollution

Nicole E. Choquette^{1,2} | Funda Ogut^{3,4} | Timothy M. Wertin¹ | Christopher M. Montes^{1,2} | Crystal A. Sorgini^{1,5} | Alison M. Morse^{3,4} | Patrick J. Brown^{1,5} | Andrew D. B. Leakey^{1,2,5} | Lauren M. McIntyre^{3,4} | Elizabeth A. Ainsworth^{1,2,5,6}

¹Carl R. Woese Institute for Genomic Biology, University of Illinois at Urbana-Champaign, Urbana, Illinois

²Department of Plant Biology, University of Illinois at Urbana-Champaign, Urbana, Illinois

³Department of Molecular Genetics and Microbiology, University of Florida, Gainesville, Florida

⁴Genetics Institute, University of Florida, Gainesville, Florida

⁵Department of Crop Sciences, University of Illinois at Urbana-Champaign, Urbana, Illinois

⁶USDA ARS Global Change and Photosynthesis Research Unit, Urbana, Illinois

Correspondence

Elizabeth A. Ainsworth, 1201 W. Gregory Drive, 147 ERML, Urbana, IL 61801.
Email: lisa.ainsworth@ars.usda.gov

Present address

Funda Ogut, Department of Forest Engineering, Artvin Coruh University, Artvin, Turkey

Funding information

National Science Foundation, Grant/Award Number: PGR-1238030

Abstract

Ozone is the most damaging air pollutant to crops, currently reducing Midwest US maize production by up to 10%, yet there has been very little effort to adapt germplasm for ozone tolerance. Ozone enters plants through stomata, reacts to form reactive oxygen species in the apoplast and ultimately decreases photosynthetic C gain. In this study, 10 diverse inbred parents were crossed in a half-diallel design to create 45 F₁ hybrids, which were tested for ozone response in the field using free air concentration enrichment (FACE). Ozone stress increased the heritability of photosynthetic traits and altered genetic correlations among traits. Hybrids from parents Hp301 and NC338 showed greater sensitivity to ozone stress, and disrupted relationships among photosynthetic traits. The physiological responses underlying sensitivity to ozone differed in hybrids from the two parents, suggesting multiple mechanisms of response to oxidative stress. FACE technology was essential to this evaluation because genetic variation in photosynthesis under elevated ozone was not predictable based on performance at ambient ozone. These findings suggest that selection under elevated ozone is needed to identify deleterious alleles in the world's largest commodity crop.

KEYWORDS

air pollution, FACE, global climate change, heritability, maize, ozone (O₃), photosynthesis

1 | INTRODUCTION

Developing crops that can be more productive under stressful growing conditions is a high priority for agriculture today, and will be increasingly necessary if we are to avoid production losses to

climate change (Challinor et al., 2014; Lesk, Rowhani, & Ramankutty, 2016; Lobell et al., 2014). Traditionally, field trials under more extreme environmental conditions than are typical for major crop growing regions have been used to test germplasm developed by breeding or biotechnology. However, some stressors—such as

Nicole E. Choquette, Funda Ogut and Timothy M. Wertin share co-first authorship.

This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

Published 2019. This article is a U.S. Government work and is in the public domain in the USA. Global Change Biology published by John Wiley & Sons Ltd.

ozone (O_3) pollution—are too heterogeneous and unpredictable in time or space to make this approach feasible (Ainsworth, Rogers, & Leakey, 2008). In addition, future climate change will result in growing environments with elevated $[CO_2]$ and temperature for which there is no present-day analogue (Battisti & Naylor, 2009; Leakey & Lau, 2012). Controlled environment growth facilities can provide valuable information on genetic variation in crop responses to stress treatments and the mechanisms underlying genetic variation (Brosché et al., 2010; Burton, Burkey, Carter, Orf, & Cregan, 2016; Frei, Tanaka, & Wissuwa, 2008; Ueda, Siddique, & Frei, 2015), but results of such controlled environment experimentation do not always translate into improved performance under production conditions in the field (Ainsworth, Beier, et al., 2008; Araus & Cairns, 2014; McKersie, Bowley, & Jones, 1999; Passioura, 2012). Free-air concentration enrichment (FACE) was developed to expose crops under field conditions to elevated concentrations of atmospheric pollutants over the entire growing season, with little or no perturbation to other aspects of the environment (Long, Ainsworth, Rogers, & Ort, 2004). But, most FACE experiments have tested a limited number of genotypes, at any given time (Betzberger et al., 2010; Markelz, Strellner, & Leakey, 2011; Wang et al., 2014). Investigation of many genotypes and structured populations is needed to understand the heritability of traits in altered atmospheric environments and ultimately to identify genomic regions and genes associated with O_3 tolerance.

Tropospheric O_3 is a dynamic, short-lived air pollutant that is estimated to cause annual losses of ~10% to US maize yields with crop losses of \$7.2 billion (McGrath et al., 2015). However, crop yield losses to O_3 pollution are not widely recognized by farmers. And, breeding or biotechnology for tolerance to O_3 stress has not been a major target for seed companies (Ainsworth, 2017). Ozone is formed as a secondary pollutant from nitrogen oxides (NOx) and volatile organic compounds, and recent analyses suggest that progress towards reducing NOx in the United States has slowed considerably, thus increasing the risks of O_3 pollution (Jiang et al., 2018). Ozone diffuses through stomatal pores on leaf surfaces and reacts to form reactive oxygen species (ROS) in the apoplast. When ROS exceed the antioxidant-quenching capacity of the apoplast, they cause oxidative stress within cells that accelerates senescence and impairs photosynthesis, ultimately reducing plant productivity and crop yields (Ainsworth, Yendrek, Sitch, Collins, & Emberson, 2012; Kangasjärvi, Jaspers, & Kollist, 2005). Maize, like many other crop species, is sensitive to O_3 damage, and shows accelerated loss of photosynthetic capacity with continued exposure to the air pollutant (Fiscus, Brooker, & Burkey, 2005; Yendrek, Erice, et al., 2017). Additionally, stomatal closure can be negatively impacted by O_3 stress, leading to excessive water loss under drought stress (Wang et al., 2014; Wilkinson & Davies, 2010). Maintenance of high photosynthetic CO_2 assimilation without excess stomatal conductance is an important phenotype for increasing O_3 tolerance (Ainsworth, 2017; Emberson et al., 2018). Furthermore, enhancing photosynthetic CO_2 assimilation and water use efficiency (ratio of photosynthetic CO_2 assimilation to water loss by transpiration) are widely

recognized to be key targets for crop improvement at a time when potential for further gains in harvest index and planting density may be limited (Leakey et al., 2019; Long, Marshall-Colon, & Zhu, 2015; von Caemmerer & Furbank, 2016). Despite the importance of maize for food, fuel and animal feed, little is known about the extent or mechanisms of genetic variation in the sensitivity of maize to O_3 by comparison to other crops such as soybean, wheat and rice (Betzberger et al., 2012; Burton et al., 2016; Frei et al., 2008; Wang et al., 2014). This represents an important unexplored opportunity because maize is a highly tractable, model system for study of crop genetics (Buckler et al., 2009; Riedelsheimer et al., 2012; Schnable et al., 2009).

For physiological performance to be a target for improvement in breeding programmes, there must be underlying additive genetic variation in the traits of interest. The likely success of selection is reflected in the narrow sense heritability, that is, the proportion of phenotypic variation resulting from additive genetic variance (Falconer & Mackay, 1996; Flood, Harbinson, & Aarts, 2011). Previous studies of maize have estimated relatively high heritability for traits related to photosynthetic capacity (Cai et al., 2012; Crosbie, Mock, & Pearce, 1977; Lu et al., 2011; Pelleschi et al., 2006; Prado et al., 2017; Wang et al., 2013; Ziyomo & Bernardo, 2013) and indicated that variance in photosynthetic traits is mostly additive (Crosbie et al., 1977). But, the heritability of photosynthetic traits in crops is reportedly lower under stress conditions (Edwards, Ewers, McClung, Lou, & Weinig, 2012; Pelleschi et al., 2006). Prior studies on leaf-level responses to O_3 in fescue, potato and sweetcorn found that additive effects (GCA), not dominance effects (SCA), were significant and involved in O_3 tolerance (De Vos, Hill, Pell, & Cole, 1982; Johnston, Haaland, & Dickens, 1983; Schraudner, Langebartels, & Sandermann, 1997). These studies fumigated crops with very high concentrations of O_3 for hours to days, which elicited acute stress responses that are known to be fundamentally distinct from responses to season-long, moderate O_3 concentrations that drive yield loss in farmer's fields (Ainsworth et al., 2012; Chen, Frank, & Long, 2009; Schraudner et al., 1997). Uncertainty regarding the extent to which photosynthetic traits associated with O_3 tolerance are heritable is compounded by the need to know if there are strong genetic correlations across environments (Falconer, 1952). In other words, if there is a substantial genotype \times environment interaction acting on photosynthetic traits, then selection for crop genotypes that can tolerate elevated O_3 pollution would not be successful under standard growing conditions. Alternatively, the absence of genotype \times environment interaction would suggest that past selection for highly productive genotypes would likely have incidentally selected for tolerance to O_3 pollution as well. In addition, genetic correlations among traits are useful in assessing how many independent traits need to be evaluated for a successful selection index to be developed.

Proof-of-concept is needed to demonstrate the use of FACE experimentation to estimate the heritability of photosynthetic traits and the degree to which elevated O_3 affects heritability in a farm-field setting (Frei, 2015). Therefore, in this study, we used a half-diallel mating design to test for (a) the effects of elevated O_3 on

not raining, when leaves were dry and when wind speed was greater than 0.5 m/s (Yendrek, Erice, et al., 2017; Yendrek, Tomaz, et al., 2017). Based on 1 min average O₃ concentration collected in each ring throughout the season, the fumigation was within 10% of the 100 nl/L target for 62% of the time and within 20% of the target concentration for 87% of the time in 2016. In 2017, fumigation was within 10% of the target for 59% of the time and within 20% of the target for 84% of the time. Other meteorological conditions were measured with an on-site weather station (Figure S1).

2.1 | Field site and experimental treatments

The O₃ treatment was applied at a target set point of 100 nL/L from 10:00 to 18:00 throughout the growing season when it was

On July 5–9, 2016 and June 28–July 1, 2017 leaf gas exchange was measured. In 2016, the eighth leaf was measured (approximately the third or fourth fully expanded leaf), while in 2017 the third fully collared leaf was measured. Prior to dawn, leaves from two plants per genotype from one ambient and one elevated O₃ plot were labelled and cut, providing ~208 leaves for measurements per day. Leaves were immediately placed in a bucket with the cut surface submerged in water, transported to a laboratory, recut under water and placed in 50 ml tubes filled with water. Leaves were stored in dim light (<100 μmol m⁻² s⁻¹). Approximately 1 hr before the measurement of leaf gas exchange, leaves were placed in a growth chamber with the following conditions: 1,800 photosynthetic photon flux density (PPFD, μmol m⁻² s⁻¹) at the leaf surface, 18°C air temperature (25°C leaf temperature) and relative humidity of ~90%. After




FIGURE 1 (a) Half-diallel design of 45 F_1 hybrids tested for response to elevated O_3 . (b) Aerial image of an elevated O_3 ring with yellow indicating the pipes from which air enriched with O_3 is released into the wind. (c) The spatial layout of a ring with different colour shading indicating the five sub-blocks within the ring. Nine genotypes were assigned to a 'set' and a common genotype (B73 \times Mo17) was replicated in each set as a spatial check (hashed boxes). Sets were randomly assigned to different sub-blocks in each of the four replicate ambient and elevated O_3 rings

approximately 1 hr of acclimation, leaf gas exchange was measured with a set of portable gas exchange systems (LI-6400; LI-COR Biosciences, Lincoln, NE, USA) with 2 × 3 red/blue LED cuvettes (LI-6400-02B; LI-COR Biosciences). Light-saturated gas exchange was measured at 1,800 PPFD ($\mu\text{mol m}^{-2} \text{s}^{-1}$), 25°C (leaf temperature) and 60% relative humidity.

Gas exchange systems were run simultaneously, and leaf mean flux of net CO_2 assimilation (A), stomatal conductance (g_s) and the ratio of the leaf intercellular concentration of CO_2 to atmospheric CO_2 ($c_i:c_a$) were measured or calculated every 4 s for 4 min (Figure S2). Analysis was done on the average values of A , g_s and $c_i:c_a$ over the last minute of measurement. Instantaneous water use efficiency ($i\text{WUE} = A/g_s$) was calculated from A and g_s . Following gas exchange measurements, leaf reflectance was measured from the adaxial surface of the leaf using a full-range spectroradiometer (ASD FieldSpec 4 Standard Res, Analytical Spectral Devices) following a standard protocol (Yendrek, Erice, et al., 2017; Yendrek, Tomaz, et al., 2017). Six reflectance measurements were collected and a splice correction was applied to the spectra to ensure continuous data across detectors (Serbin, Singh, McNeil, Kingdon, & Townsend, 2014). Those six spectra from each leaf were then averaged and data were interpolated to provide 1 nm bandwidths. Further quality control was applied using the FieldSpectra package in R (Serbin et al., 2014). Partial least squares regression models were then applied to the spectra to estimate the CO_2 -saturated rate of A (V_{maxm}) and leaf chlorophyll content (Chl_m ; Yendrek, Erice, et al., 2017; Yendrek, Tomaz, et al., 2017).

2.3 | Data quality control

Measurements of gas exchange over time were examined for data quality and conformation to basic assumptions. If mean g_s or c_i for the last minute of the measurement period were negative, then data were not used (Figure S2). Individual leaf measurements for the same genotype and treatment were evaluated for outliers and influential observations. If an individual measurement strongly influenced the stability of the data, the observation was not used for further statistical analysis.

2.4 | Statistical analysis

B73 × Mo17 was planted within each sub-block of the ring (Figure 1), and used as a covariate in the model (see Supplemental Text for additional statistical tests). As the majority of traits showed evidence for interactions between additive genetic variance and an interaction between additive genetic variance and O_3 treatment, the ambient and elevated O_3 environments were modelled separately to estimate heritability:

$$y_{ijkm} = \mu_i + C_{ijk} + G_m + \varepsilon_{ijkm}.$$

y_{ijkm} is the observed trait value for the i th treatment (ambient or elevated O_3) in the k th sub-block of the j th ring-pair, belonging to the cross from the m th mother where μ_i is the mean for the i th

treatment. C_{ijk} is the trait value of the B73 × Mo17 check plot in the i th treatment, j th ring-pair and k th sub-block. G_m is the random GCA effect (parameterized as a set of indicator variables for the maternal parent). The variance component for GCA was estimated using a Toeplitz(1) variance structure (Wayne et al., 2007).

Narrow sense heritability and standard error (SE) were estimated for each trait in ambient and elevated O_3 conditions separately as:

$$h_n^2 = \frac{2 * \sigma_{\text{gca}}^2}{2 * \sigma_{\text{gca}}^2 + \sigma_e^2},$$

where σ_{gca}^2 estimated variance of GCA effect and σ_e^2 is residual variance. The SE of heritability estimates was obtained with the Delta method (Wayne et al., 2007).

Genetic correlations between environments for the same trait ($r_{\text{g_amb_oz}}$) were calculated using a MANOVA based on the above model where the two dependent variables were the genotypic mean values in each of the two environments:

$$Y_i = \begin{bmatrix} w_{\text{amb}} \\ w_{\text{oz}} \end{bmatrix}, \quad \varepsilon_i = \begin{bmatrix} \varepsilon_{\text{amb}} \\ \varepsilon_{\text{oz}} \end{bmatrix}$$

$$Z_i \sim \left(0, \begin{bmatrix} \sigma_{\text{gca_amb}}^2 & \sigma_{\text{gca_amb_oz}} \\ \sigma_{\text{gca_amb_oz}} & \sigma_{\text{gca_oz}}^2 \end{bmatrix} \right).$$

The genetic correlation between environments for the same trait was calculated from estimates in the above model as:

$$r_{\text{g_amb_oz}} = \frac{\sigma_{\text{gca_amb_oz}}}{\sqrt{\sigma_{\text{gca_amb}}^2 * \sigma_{\text{gca_oz}}^2}},$$

where $\sigma_{\text{gca_amb_oz}}$ is the genetic covariance between environments for a given trait and $\sigma_{\text{gca_amb}}^2$ and $\sigma_{\text{gca_oz}}^2$ are the additive genetic variances for the same trait in ambient and elevated O_3 respectively.

Genetic correlations between traits within an O_3 treatment ($r_{\text{g_trait1_trait2}}$) were calculated based on the above model using a MANOVA approach where a pair of traits was considered within the same environment:

$$Y_i = \begin{bmatrix} w_{\text{trait1_oz}} \\ w_{\text{trait2_oz}} \end{bmatrix}, \quad \varepsilon_i = \begin{bmatrix} \varepsilon_{\text{trait1_oz}} \\ \varepsilon_{\text{trait2_oz}} \end{bmatrix}$$

$$\gamma_i = \begin{bmatrix} \gamma_{\text{trait1_oz}} \\ \gamma_{\text{trait2_oz}} \end{bmatrix} \sim \left(0, \begin{bmatrix} \sigma_{\text{gca_trait1_oz}}^2 & \sigma_{\text{gca_trait1_oz_trait2_oz}} \\ \sigma_{\text{gca_trait1_oz_trait2_oz}} & \sigma_{\text{gca_trait2_oz}}^2 \end{bmatrix} \right).$$

Genetic correlations between traits for each environment were calculated as:

$$r_{\text{g_trait1_trait2_oz}} = \frac{\sigma_{\text{gca_trait1_oz_trait2_oz}}}{\sqrt{\sigma_{\text{gca_trait1_oz}}^2 * \sigma_{\text{gca_trait2_oz}}^2}},$$

where $\sigma_{\text{gca_trait1_oz_trait2_oz}}$ is the additive genetic covariance between traits for the same environment, and $\sigma_{\text{gca_trait1_oz}}^2$ and $\sigma_{\text{gca_trait2_oz}}^2$ are the additive genetic variances for trait1 and trait2.

Phenotypic correlations within an O_3 treatment ($r_{p_trait1_trait2}$) were calculated as:

$$r_{p_trait1_trait2} = \frac{\sigma_{p_trait1_oz_trait2_oz}}{\sqrt{\sigma_{p_trait1_oz}^2 * \sigma_{p_trait2_oz}^2}},$$

where $\sigma_{p_trait1_oz_trait2_oz} = \sigma_{gca_trait1_oz_trait2_oz} + \sigma_{e_trait1_oz_trait2_oz}$. All calculations were performed using code adapted from Holland (2006).

Visual examination of the results led to the hypothesis that NC338 and HP301 were responsible for the shift in genetic correlations. Thus, the correlation analyses described above were repeated for the set of F_1 hybrids that did not include hybrids NC338 or HP301 as either parent.

3 | RESULTS

3.1 | Gas exchange response to elevated O_3

Photosynthetic traits of 45 F_1 hybrids comprising a half-diallel population were measured in ambient and elevated O_3 in 2016 and 2017. Measurements were taken on mature leaves approximately midway through vegetative development of the crop in both years. Rainfall was

lower in 2017, leading to greater drought stress than in 2016 (Figure S2), and lower A and g_s (Figure 2; Figure S3). In both years, averaging across hybrids, elevated O_3 reduced A , by ~14% (Figure 2), and g_s by 6% in 2016 and by 14% in 2017 (Figure S3).

3.2 | Narrow sense heritability increases with O_3 environment

Analysis of the half-diallel maize population grown showed that additive genetic variance explained the variation in photosynthetic and reflectance traits (Table S1). There was little evidence for dominance variance (Table S1; comparison of models 2 and 3; Table S2). In addition, there was strong evidence for interactions between genetic variation and environment ($G \times E$; Table S1; comparison of models 2 and 3), indicating that the genetic architecture of most traits was different in the two environments. Therefore, heritability was calculated separately in ambient and elevated O_3 . Narrow sense heritability (h_n^2) estimates of photosynthetic traits differed between ambient and elevated O_3 in 2016 and 2017 (Table 1). Interestingly, h_n^2 in elevated O_3 was greater than in ambient O_3 for many photosynthetic traits (Table 1), indicating the presence of additional additive genetic variation in elevated O_3 . Estimated h_n^2 of A was greater than h_n^2 of other




FIGURE 2 Measurements of net carbon assimilation (A) measured in the F_1 maize hybrids. Measurements were made during the summer of 2016 (a) and 2017 (b) on hybrids grown in ambient O_3 (white) and elevated O_3 (black). Data are ordered based on A measured at ambient O_3 . Error bars represent 1 SD

| | 2016 | | 2017 | |
|------------------|------------------------|-------------------------|------------------------|-------------------------|
| | Ambient O ₃ | Elevated O ₃ | Ambient O ₃ | Elevated O ₃ |
| A | 0.222 | 0.678 | 0.315 | 0.499 |
| g_s | 0.178 | 0.380 | 0.278 | 0.202 |
| iWUE | 0.127 | 0.182 | 0.212 | 0.203 |
| $c_i:c_a$ | 0.103 | 0.182 | 0.197 | 0.237 |
| V_{max} | 0.125 | 0.236 | 0.197 | 0.258 |
| Chl _m | 0.424 | 0.283 | 0.171 | 0.276 |

TABLE 1 Narrow sense heritability (h_n^2) for photosynthetic traits measured in 2016 and 2017 in ambient and elevated [O₃]




FIGURE 3 Genetic and phenotypic correlations among photosynthetic traits. Genetic correlations (a, c) and phenotypic correlations (b, d) among photosynthetic traits in ambient and elevated O₃ calculated in 2016 and 2017. Top panels include all hybrids (a, b). Bottom panels show correlations after hybrids with parents Hp301 or NC338 were removed from the analysis (b, c)

photosynthetic traits (Table 1), and was of similar magnitude to previous estimates in maize (Crosbie et al., 1977; Pelleschi et al., 2006).

3.3 | Elevated O₃ alters genetic and phenotypic correlations among photosynthetic traits

Genetic correlations (R_g) are important to define the shared genetic components between traits and phenotypic correlations (R_p)

measure the consistency of performance between traits. Both R_g and R_p between A and g_s were very strong regardless of the environment or year (Figure 3a,b, red symbols). This is unsurprising because g_s is a principle determinant of CO₂ entry into leaves, and A measures CO₂ fixation. R_g between g_s and iWUE were strongly negative in both 2016 and 2017, indicating that selection for high g_s in maize would result in low iWUE (Figure 3, grey symbols). There was a positive R_g between traits estimated from leaf reflectance




FIGURE 4 The relationship between stomatal conductance (g_s) and the ratio of the leaf intercellular concentration of CO_2 to atmospheric CO_2 ($c_i:c_a$) in maize hybrids grown at ambient (circles) and elevated O_3 (triangles) in 2016 (a) and 2017 (b). Blue symbols represent hybrids with Hp301 as a parent and orange symbols hybrids with NC338 as a parent




FIGURE 5 Per cent change of photosynthetic traits at elevated O_3 . Blue circles represent hybrids with Hp301 as the male or female parent and orange circles are hybrids with NC338 as the male or female parent

spectra, Chl_m and V_{\max} (Figure 3a, olive symbols) in both ambient and elevated O_3 , but R_g between A and V_{\max} or Chl_m was not strong or consistent (Figure 3a, pink symbols). It was notable that R_g between A and $c_i:c_a$ (Figure 3a, green symbols) and g_s and $c_i:c_a$ (Figure 3a, cyan symbols) differed in ambient and elevated O_3 . R_g between g_s and $c_i:c_a$ was strong and positive in ambient O_3 , but not in elevated O_3 (Figure 3a, cyan symbols), and R_g between A and $c_i:c_a$ was positive in ambient O_3 and negative in elevated O_3 (Figure 3a, green symbols). These same general trends were also observed for R_p (Figure 3b).

3.4 | Identification of O_3 -sensitive maize lines

The correlation between g_s and $c_i:c_a$ was further explored based on the differences in R_g between traits in ambient and elevated O_3 . Across the diverse F_1 hybrids, there was a strong, linear relationship between g_s and $c_i:c_a$ consistent among all lines in ambient O_3 (Figure 4, circles). However, growth at elevated O_3 altered the relationship between g_s and $c_i:c_a$ in F_1 crosses with parents NC338 (Figure 4, orange triangles) and Hp301 (Figure 4, blue triangles) in both 2016 and 2017. Notably, alleles from Hp301 and NC338 appear to confer sensitivity to elevated O_3 in different ways. Lower A at elevated O_3 in hybrids with Hp301 as a parent was linked to lower g_s at elevated O_3 without a change in $c_i:c_a$ (Figure 4). Meanwhile, lower A at elevated O_3 in hybrid with NC338 as a parent was linked to greater $c_i:c_a$ without a change in g_s (Figure 4). F_1 hybrids with these two parents also showed the greatest per cent decrease in A and g_s at elevated O_3 in 2016 and 2017 (Figure 5a,b). When R_g and R_p were estimated without hybrids containing NC338 or Hp301 (Figure 3c,d), then estimates were similar in both ambient and elevated O_3 . This further indicates that F_1 crosses with Hp301 and NC338 are sensitive to O_3 and drive the differences in R_g and R_p observed between A and $c_i:c_a$ and g_s and $c_i:c_a$.

4 | DISCUSSION

This 2-year study enabled investigation of genetic variation in photosynthetic traits as well as the consistency of the response of photosynthetic traits to elevated O_3 across growing seasons. Photosynthesis has been suggested as one of the most important processes to improve in order to combat stagnating crop yields and improve future food security (Long et al., 2015). Transgenic approaches to improve C4 photosynthesis have been suggested (von Caemmerer & Furbank, 2016), but exploiting standing genetic variation in photosynthetic traits is also a path to crop improvement that does not involve transgenic technology (Cañas et al., 2017). Here, we showed that maize hybrids have significant additive genetic variation in A in both ambient and elevated O_3 (Figure 2), and the per cent change in A at elevated O_3 was consistent in both years (Figure 5a). By using a half-diallel mating design comprised of diverse maize lines, we estimated genetic (R_g) and phenotypic (R_p) correlations in both environments and demonstrated that growth at elevated O_3 alters the R_g and R_p between photosynthetic traits. Growth at

elevated O_3 changes the relationship between g_s and $c_i:c_a$, primarily driven by genotypes with Hp301 and NC338. Furthermore, the mechanisms of sensitivity to elevated O_3 appear to differ between genotypes Hp301 and NC338. Overall, we have demonstrated the utility of FACE experiments for screening structured populations to estimate heritability of phenotypic traits and how those change with O_3 pollution. This capacity and information is critical for any future efforts to breed for O_3 tolerance.

Previous studies of O_3 -induced leaf damage in fescue, tobacco, maize, potato and *Plantago major* also showed little evidence of dominance variance (Aycock, 1972; Cameron, 1975; Campbell, Devine, & Howelp, 1977; De Vos et al., 1982; Huang, Aycock, & Mulchi, 1975; Johnston et al., 1983; Whitfield, Davison, & Ashenden, 1997), which is consistent with our findings. Lack of dominance variance indicates that the narrow sense heritability (h_n^2) of net carbon assimilation (A) in elevated O_3 is mainly additive and can be attributed to additive genetic factors inherited from the parents. The observed increase in h_n^2 of A under O_3 stress is notable. A prior study investigating O_3 sensitivity in Black Cherry (*Prunus serotina* Ehrh.) also found that heritability of the foliar injury increased with exposure to greater treatment concentrations of elevated O_3 (Lee, Steiner, Zhang, & Skelly, 2002). In contrast, studies that investigated other abiotic stresses such as drought stress often reported that h_n^2 was lower under stress conditions. For example, drought stress decreased h_n^2 of A from 0.61 to 0.33 in *Brassica rapa* (Edwards et al., 2012) and from 0.45 to 0.37 in maize (Pelleschi et al., 2006). Greater h_n^2 in elevated O_3 as revealed in this study suggests that there is potential for improvement. Furthermore, selection for A under standard growing conditions of the Midwest in the United States would fail to select against deleterious alleles that confer sensitivity to elevated O_3 (Gibson & Dworkin, 2004).

Genetic correlations (R_g) among traits imply that the same genes are acting on multiple traits. If the correlation is high enough, it is possible that selection can be performed on one trait, with the second trait also improving due to shared loci between the traits. We tested R_g among photosynthetic traits measured with gas exchange and traits estimated from leaf hyperspectral reflectance (Yendrek, Erice, et al., 2017; Yendrek, Tomaz, et al., 2017), which has been suggested as a promising high-throughput approach for phenotyping (Araus & Cairns, 2014; Furbank & Tester, 2011). Although R_g between V_{\max} and Chl_m estimated from hyperspectral reflectance was strong and positive, R_g between gas exchange traits and remotely sensed traits was not strong in our experiments (Figure 3a). This could be because chlorophyll content and maximum photosynthetic capacity were not the primary limitations to gas exchange at the times of our measurement. In 2017, reflectance was measured in the field on intact plants, which may also lead to lower correlations. We did observe strong, positive correlations between A and g_s , which has been reported previously for other species under variable environmental conditions (Manzaneda, Rey, Anderson, Raskin, & Mitchell, 2016; Pelleschi et al., 2006). We also identified a fundamental shift in R_g between

g_s and $c_i:c_a$ in ambient and elevated O_3 (Figure 3a). $c_i:c_a$ provides information on the balance between resistance for CO_2 diffusion into the leaf and the biochemical capacity for CO_2 fixation in the mesophyll. Altered R_g in elevated O_3 suggests the involvement of additional genetic factors in controlling the phenotype under O_3 stress.

Further analysis of the relationship between g_s and $c_i:c_a$ identified that hybrids with NC338 and Hp301 were more sensitive to O_3 stress (Figure 4). When R_g and R_p were calculated without hybrids containing Hp301 and NC338, the correlations become similar in ambient and elevated O_3 (Figure 3c,d), suggesting that alleles from Hp301 and NC338 were responsible for the change in genetic architecture of photosynthetic traits in elevated O_3 . These alleles from Hp301 and NC338 demonstrate different mechanisms of sensitivity to elevated O_3 , with Hp301 crosses linked to lower g_s at elevated O_3 with no change in $c_i:c_a$ and NC338 hybrids showing greater $c_i:c_a$ without a change in g_s (Figure 5). This implies that stomatal function was disrupted in Hp301 lines, which is consistent with the prior evidence that O_3 stress alters stomatal signalling and closure (Wilkinson & Davies, 2010). Meanwhile, our findings suggest that alterations in mesophyll conductance (g_m) or perhaps decreases in photosynthetic capacity drove sensitivity to elevated O_3 in hybrids with NC338 as a parent. This is consistent with greater average reductions in V_{maxm} and Chl_m in hybrids containing NC338 as a parent (Figure 5). A study of Siebold's beech (*Fagus crenata*) seedlings found a significant increase in c_i and a substantial decrease in g_m with increasing O_3 stress, suggesting that lower g_m hindered CO_2 transfer to the site of carbon fixation (Watanabe et al., 2018). Similarly, a study on snap peas found the genotype sensitive to elevated O_3 had a significant decrease in g_m (Flowers, Fiscus, Burkey, Booker, & Dubios, 2007). Photosynthetic processes may be sensitive to O_3 (Fiscus et al., 2005) and reductions in PEPc activity and Rubisco activity were reported at elevated O_3 in young maize leaves (Leitao, Bethenod, & Biolley, 2007). Such reductions in either g_m or photosynthetic enzyme activity if decoupled from g_s could increase c_a , as observed in the sensitive NC338 hybrids here.

Global mean O_3 pollution increased by 8.9% from 1990 to 2013 (Brauer et al., 2016) causing a significant, negative impact on crop production in North America and Europe worldwide (Mills et al., 2018). Identifying genetic variation in photosynthetic traits and tolerant genotypes to elevated O_3 is a crucial step towards breeding for O_3 tolerance (Ainsworth, 2017). For breeding to be straightforward, genetic variation should be directly available to selection, and the genetic contribution to phenotypic variation should be additive. This first investigation of a half-diallel population of maize exposed to elevated O_3 in the field showed not only that additive genetic variation is a large component of phenotypic variation in photosynthetic traits, but also that O_3 stress increased the heritability of photosynthetic traits, indicating additional loci responding to O_3 stress. In addition, the genetic correlation among photosynthetic traits changed between environments. The F_1 hybrids from parents Hp301 and NC338 were particularly sensitive to O_3 and showed disrupted relationships between A or

g_s and $c_i:c_a$. The physiological mechanism underlying this sensitivity was distinct between the two parent lines, suggesting that multiple sources of sensitivity to oxidative stress from air pollution might need to be selected against to optimize maize production. This implies that past selection of maize under ambient O_3 did not select for alleles that confer tolerance to elevated O_3 pollution. Further selection under elevated O_3 concentrations might purge deleterious alleles in the world's most important commodity crop. More broadly, this work demonstrates the capability of FACE technology to be used for field experimentation on diverse populations of major commodity crops to address needs for adaptation to pollution and climatic change that cannot be easily accomplished by other existing approaches.

ACKNOWLEDGEMENTS

This work was supported by a grant from the NSF Plant Genome Research Program (PGR-1238030). We thank Brad Dalsing, Chad Lance, Aidan McMahon and Jesse McGrath for operating fumigation at SoyFACE. We thank Chris Moller, Ben Thompson, John Ferguson, Charlie Burroughs, Lorena Rios-Acosta and Jessica Wedow for help with sampling, photosynthetic and reflectance measurements. We thank Jim Berry, Aya Bridgeland, Cecilia Pigozzi, Savannah Pflugmacher, Noah Mitchell and Emma Diers for their help in maintaining the research plots. We thank Linda Young for assistance with the spatial design of the experiment. Any opinions, findings and conclusions or recommendations expressed in this publication are those of the author(s) and do not necessarily reflect the views of the U.S. Department of Agriculture. Mention of trade names or commercial products in this publication is solely for the purpose of providing specific information and does not imply recommendation or endorsement by the U.S. Department of Agriculture. USDA is an equal opportunity provider and employer.

ORCID

Andrew D. B. Leakey  <https://orcid.org/0000-0001-6251-024X>

Lauren M. McIntyre  <https://orcid.org/0000-0002-0077-3359>

Elizabeth A. Ainsworth  <https://orcid.org/0000-0002-3199-8999>

REFERENCES

- Ainsworth, E. A. (2017). Understanding and improving global crop response to ozone pollution. *Plant Journal*, 90(5), 886–897. <https://doi.org/10.1111/tpj.13298>
- Ainsworth, E. A., Beier, C., Calfapietra, C., Reinhart, C., Durand-Tardif, M., Graham, F. D., ... White, J. W. (2008). Next generation of elevated $[CO_2]$ experiments with crops: A critical investment for feeding the future world. *Plant Cell & Environment*, 31(9), 1317–1324.
- Ainsworth, E. A., Rogers, A., & Leakey, A. D. B. (2008). Targets for crop biotechnology in a future high- CO_2 and high- O_3 world. *Plant Physiology*, 147(1), 13–19.

- Ainsworth, E. A., Yendrek, C. R., Sitch, S., Collins, W. J., & Emberson, L. D. (2012). The effects of tropospheric ozone on net primary productivity and implications for climate change. *Annual Review of Plant Biology*, 63(1), 637–661. <https://doi.org/10.1146/annurev-arplant-042110-103829>
- Araus, J. L., & Cairns, J. E. (2014). Field high-throughput phenotyping: The new crop breeding frontier. *Trends in Plant Science*, 19(1), 52–61. <https://doi.org/10.1016/j.tplants.2013.09.008>
- Aycock, M. K. (1972). Combining ability estimates for weather fleck in *Nicotiana tabacum* L. *Crop Science*, 12(5), 672–674.
- Battisti, D. S., & Naylor, R. L. (2009). Historical warnings of future food insecurity with unprecedented seasonal heat. *Science*, 323(5911), 240–244. <https://doi.org/10.1126/science.1164363>
- Betzberger, A. M., Gillespie, K. M., McGrath, J. M., Koester, R. P., Nelson, R. L., & Ainsworth, E. A. (2010). Effects of chronic elevated ozone concentration on antioxidant capacity, photosynthesis and seed yield of 10 soybean cultivars. *Plant Cell Environment*, 33(9), 1569–1581. <https://doi.org/10.1111/j.1365-3040.2010.02165.x>
- Betzberger, A. M., Yendrek, C. R., Sun, J., Leisner, C. P., Nelson, R. L., Ort, D. R., & Ainsworth, E. A. (2012). Ozone exposure response for U.S. soybean cultivars: Linear reductions in photosynthetic potential, biomass, and yield. *Plant Physiology*, 160(4), 1827–1839. <https://doi.org/10.1104/pp.112.205591>
- Brauer, M., Freedman, G., Frostad, J., Van Donkelaar, A., Martin, R. V., Dentener, F., ... Cohen, A. (2016). Ambient air pollution exposure estimation for the global burden of disease 2013. *Environmental Science & Technology*, 50(1), 79–88. <https://doi.org/10.1021/acs.est.5b03709>
- Brosché, M., Merilo, E., Mayer, F., Pechter, P., Puzörjova, I., Brader, G., ... Kollist, H. (2010). Natural variation in ozone sensitivity among *Arabidopsis thaliana* accessions and its relation to stomatal conductance. *Plant Cell and Environment*, 33(6), 914–925. <https://doi.org/10.1111/j.1365-3040.2010.02116.x>
- Buckler, E. S., Holland, J. B., Bradbury, P. J., Acharya, C. B., Brown, P. J., Browne, C., ... McMullen, M. D. (2009). The genetic architecture of maize flowering time. *Science*, 325(5941), 714–718. <https://doi.org/10.1126/science.1174276>
- Burton, A. L., Burkey, K. O., Carter, T. E. Jr, Orf, J., & Cregan, P. (2016). Phenotypic variation and identification of quantitative trait loci for ozone tolerance in a Fiskeby III × Mandarin (Ottawa) soybean population. *Theoretical and Applied Genetics*, 129(6), 1113–1125. <https://doi.org/10.1007/s00122-016-2687-1>
- Cai, Q. S., Wang, L. L., Yao, W. H., Zhang, Y. D., Liu, L., Yu, L. J., & Fan, X. M. (2012). Diallel analysis of photosynthetic traits in maize. *Crop Science*, 52(2), 551–559. <https://doi.org/10.2135/cropsci2011.06.0333>
- Cameron, J. W. (1975). Inheritance in sweet corn for resistance to acute ozone injury. *Journal of the American Society for Horticultural Science*, 100(5), 577–657.
- Campbell, T. A., Devine, T. E., & Howelp, R. K. (1977). Diallel analysis of resistance to air pollutants in alfalfa. *Crop Science*, 17(4), 664–665. <https://doi.org/10.2135/cropsci1977.0011183x001700040049x>
- Cañas, R. A., Yesberger-nove-Cuny, Z., Simons, M., Chardon, F., Armengaud, P., Quillere, I., ... Hirel, B. (2017). Exploiting the genetic diversity of maize using a combined metabolomic, enzyme activity profiling, and metabolic approach to link leaf physiology to kernel yield. *The Plant Cell*, 29(5), 919–943. <https://doi.org/10.1105/tpc.16.00613>
- Challinor, A. J., Watson, J., Lobell, D. B., Howden, S. M., Smith, D. R., & Chhetri, N. (2014). A meta-analysis of crop yield under climate change and adaptation. *Nature Climate Change*, 4(4), 287–291. <https://doi.org/10.1038/nclimate2153>
- Chen, C. P., Frank, T. D., & Long, S. P. (2009). Is a short, sharp shock equivalent to long-term punishment? Contrasting the spatial pattern of acute and chronic ozone damage to soybean leaves via chlorophyll fluorescence imaging. *Plant Cell and Environment*, 32(4), 327–335. <https://doi.org/10.1111/j.1365-3040.2008.01923.x>
- Crosbie, T. M., Mock, J. J., & Pearce, R. B. (1977). Variability and selection advance for photosynthesis in Iowa stiff stalk synthetic maize population. *Crop Science*, 17(4), 511–514. <https://doi.org/10.2135/cropsci1977.0011183x001700040007x>
- De Vos, N. E., Hill, R. R., Pell, E. J., & Cole, R. H. (1982). Quantitative inheritance of ozone resistance in potato. *Crop Science*, 22(5), 992–995. <https://doi.org/10.2135/cropsci1982.0011183x002200050023x>
- Edwards, C. E., Ewers, B. E., McClung, C. R., Lou, P., & Weinig, C. (2012). Quantitative variation in water-use efficiency across water regimes and its relationship with circadian, vegetative, reproductive, and leaf gas-exchange traits. *Molecular Plant*, 5(3), 653–668. <https://doi.org/10.1093/mp/sss004>
- Emberson, L. D., Pleijel, H., Ainsworth, E. A., van den Berg, M., Ren, W., Osborne, S., ... Van Dingenen, R. (2018). Ozone effects on crops and consideration in crop models. *European Journal of Agronomy*, 100, 19–34. <https://doi.org/10.1016/j.eja.2018.06.002>
- Falconer, D. S. (1952). The problem of environment and selection. *American Naturalist*, 86(830), 293–298. <https://doi.org/10.1086/281736>
- Falconer, D. S., & Mackay, T. F. C. (1996). *Introduction to quantitative genetics* (4th ed.). Essex, UK: Prentice Hall.
- Fiscus, E. L., Brooker, F. L., & Burkey, K. O. (2005). Crop responses to ozone: Uptake, modes of action, carbon assimilation and partitioning. *Plant, Cell and Environment*, 28(8), 997–1011. <https://doi.org/10.1111/j.1365-3040.2005.01349.x>
- Flood, P. J., Harbinson, J., & Aarts, M. G. M. (2011). Natural genetic variation in plant photosynthesis. *Trends in Plant Science*, 16(6), 327–335. <https://doi.org/10.1016/j.tplants.2011.02.005>
- Flowers, M. D., Fiscus, E. L., Burkey, K. O., Booker, F. L., & Dubois, J. B. (2007). Photosynthesis, chlorophyll fluorescence, and yield of snap bean (*Phaseolus vulgaris* L.) genotypes differing in sensitivity to ozone. *Environmental and Experimental Botany*, 61(2), 190–198. <https://doi.org/10.1016/j.envexpbot.2007.05.009>
- Frei, M. (2015). Breeding of ozone resistant rice: Relevance, approaches and challenges. *Environmental Pollution*, 197, 144–155. <https://doi.org/10.1016/j.envpol.2014.12.011>
- Frei, M., Tanaka, J. P., & Wissuwa, M. (2008). Genotypic variation in tolerance to elevated ozone in rice: Dissection of distinct genetic factors linked to tolerance mechanisms. *Journal of Experimental Botany*, 59(13), 3741–3752. <https://doi.org/10.1093/jxb/ern222>
- Furbank, R. T., & Tester, M. (2011). Phenomics: Technologies to relieve the phenotyping bottleneck. *Trends in Plant Science*, 16(12), 635–644. <https://doi.org/10.1016/j.tplants.2011.09.005>
- Gibson, G., & Dworkin, I. (2004). Uncovering cryptic genetic variation. *Nature Reviews Genetics*, 5(9), 681–690. <https://doi.org/10.1038/nrg1426>
- Holland, J. B. (2006). Estimating genotypic correlations and their standard errors using multivariate restricted maximum likelihood estimation with SAS Proc Mixed. *Crop Science*, 46(2), 642–654. <https://doi.org/10.2135/cropsci2005.0191>
- Huang, T. R., Aycock, M. K., & Mulchi, C. L. (1975). Heterosis and combining ability estimates for air pollution damage, primarily ozone, in Maryland tobacco. *Crop Science*, 15(60), 785–789. <https://doi.org/10.2135/cropsci1975.0011183x001500060014x>
- Jiang, Z., McDonald, B. C., Worden, H., Worden, J. R., Miyazaki, K., Qu, Z., ... Boersma, K. F. (2018). Unexpected slowdown of US pollutant emission reduction in the past decade. *Proceedings of the National Academy of Sciences of the United States of America*, 115(20), 5099–5104. <https://doi.org/10.1073/pnas.1801191115>
- Johnston, W. J., Haaland, R. L., & Dickens, R. (1983). Inheritance of ozone resistance in tall fescue. *Crop Science*, 23(2), 235–236. <https://doi.org/10.2135/cropsci1983.0011183x002300020011x>
- Kangasjärvi, J., Jaspers, P., & Kollist, H. (2005). Signalling and cell death in ozone-exposed plants. *Plant, Cell and Environment*, 28(8), 1021–1036. <https://doi.org/10.1111/j.1365-3040.2005.01325.x>

- Leakey, A. D. B., Ferguson, J. N., Pignon, C. P., Wu, A., Jin, Z., Hammer, G. L., & Lobell, D. B. (2019). Water use efficiency as a constraint and target for improving the resilience and productivity of C3 and C4 crops. *Annual Review of Plant Biology*, 70, 781–808. <https://doi.org/10.1146/annurev-arplant-042817-040305>
- Leakey, A. D. B., & Lau, J. A. (2012). Evolutionary context for understanding and manipulating plant responses to past, present and future atmospheric CO_2 . *Philosophical Transactions of the Royal Society B*, 367(1588), 613–629. <https://doi.org/10.1098/rstb.2011.0248>
- Lee, J. C., Steiner, K. C., Zhang, J. W., & Skelly, J. M. (2002). Heritability of ozone sensitivity in open-pollinated families of black cherry (*Prunus serotina* Ehrh.). *Forest Science*, 48(1), 111–117. <https://doi.org/10.1093/forestscience/48.1.111>
- Leitao, L., Bethenod, O., & Biolley, J. (2007). The impact of ozone on juvenile maize (*Zea mays* L.) plant photosynthesis: Effects on vegetative biomass, pigmentation, and carboxylases (PEPc and Rubisco). *Plant Biology*, 9(4), 478–488. <https://doi.org/10.1055/s-2007-964942>
- Lesk, C., Rowhani, P., & Ramankutty, N. (2016). Influence of extreme weather disasters on global crop production. *Nature*, 529(7584), 84–87. <https://doi.org/10.1038/nature16467>
- Lobell, D. B., Roberts, M. J., Schlenker, W., Braun, N., Little, B. B., Rejesus, R. M., & Hammer, G. L. (2014). Greater sensitivity to drought accompanies maize yield increase in the U.S. Midwest. *Science*, 344(6183), 516–519. <https://doi.org/10.1126/science.1251423>
- Long, S. P., Ainsworth, E. A., Rogers, A., & Ort, D. R. (2004). Rising atmospheric carbon dioxide: Plants FACE the future. *Annual Review of Plant Biology*, 55, 591–628. <https://doi.org/10.1146/annurev.arplant.55.031903.141610>
- Long, S. P., Marshall-Colon, A., & Zhu, X.-G. (2015). Meeting the global food demand of the future by engineering crop photosynthesis and yield potential. *Cell*, 161(1), 56–66. <https://doi.org/10.1016/j.cell.2015.03.019>
- Lu, Y., Hao, Z., Xie, C., Crossa, J., Araus, J.-L., Gao, S., ... Xu, Y. (2011). Large-scale screening for maize drought resistance using multiple selection criteria evaluated under water-stressed and well-watered environments. *Field Crops Research*, 124(1), 37–45. <https://doi.org/10.1016/j.fcr.2011.06.003>
- Manzaneda, A. J., Rey, P. J., Anderson, J. T., Raskin, E., & Mitchell-Olds, T. (2016). Natural variation, differentiation and genetic tradeoffs of ecophysiological traits in response to water limitation in *Brachypodium distachyon* and its descendent allotetraploid *B. hybridum* (Poaceae). *Evolution*, 69(10), 2689–2704. <https://doi.org/10.1111/evo.12776>
- Markelz, R. J. C., Strellner, R. S., & Leakey, A. D. B. (2011). Impairment of C4 photosynthesis by drought is exacerbated by limiting nitrogen and ameliorated by elevated CO_2 in maize. *Journal of Experimental Botany*, 62(9), 3235–3246. <https://doi.org/10.1093/jxb/err056>
- McGrath, J. M., Betzelberger, A. M., Wang, S., Shook, E., Zhu, X.-G., Long, S. P., & Ainsworth, E. A. (2015). An analysis of ozone damage to historical maize and soybean yields in the United States. *Proceedings of the National Academy of Sciences of the United States of America*, 112(46), 14390–14395. <https://doi.org/10.1073/pnas.1509777112>
- McKersie, B. D., Bowley, S. R., & Jones, K. S. (1999). Winter survival of transgenic alfalfa overexpressing superoxide dismutase. *Plant Physiology*, 119(3), 839–847. <https://doi.org/10.1104/pp.119.3.839>
- Mills, G., Sharps, K., Simpson, D., Pleijel, H., Frei, M., Burkey, K., ... Agrawal, M. (2018). Closing the global ozone yield gap: Quantification and cobenefits for multistress tolerance. *Global Change Biology*, 24(10), 4869–4893. <https://doi.org/10.1111/gcb.14381>
- Passioura, J. B. (2012). Phenotyping for drought tolerance in grain crops: When is it useful to breeders? *Functional Plant Biology*, 39(10–11), 851–859. <https://doi.org/10.1071/FP12079>
- Pelleschi, S., Leonardi, A., Rocher, J. P., Cornic, G., De Vienne, D., Thévenot, C., & Prioul, J. L. (2006). Analysis of the relationships between growth, photosynthesis and carbohydrate metabolism using quantitative trait loci (QTLs) in young maize plants subjected to water deprivation. *Molecular Breeding*, 17(1), 21–39. <https://doi.org/10.1007/s11032-005-1031-2>
- Prado, S. A., Cabrera-Bosquet, L., Grau, A., Coupel-Ledru, A., Millet, E. J., Welcker, C., & Tardieu, F. (2017). Phenomics allows identification of genomic regions affecting maize stomatal conductance with conditional effects of water deficit and evaporative demand. *Plant Cell and Environment*, 41(2), 314–326. <https://doi.org/10.1111/pce.13083>
- Riedelshimer, C., Czedik-Eysenberg, A., Grider, C., Lisec, J., Technow, F., Sulpice, R., ... Melchinger, A. E. (2012). Genomic and metabolic prediction of complex heterotic traits in hybrid maize. *Nature Genetics*, 44(2), 217–220. <https://doi.org/10.1038/ng.1033>
- Schnable, P. S., Ware, D., Fulton, R. S., Stein, J. C., Wei, F., Pasternak, S., ... Wilson, R. K. (2009). The B73 maize genome: Complexity, diversity, and dynamics. *Science*, 326(5956), 1112–1115. <https://doi.org/10.1126/science.1178534>
- Schraudner, M., Langebartels, C., & Sandermann, H. (1997). Changes in the biochemical status of plant cells induced by the environmental pollutant ozone. *Physiologia Plantarum*, 100(2), 274–280. <https://doi.org/10.1111/j.1399-3054.1997.tb04783.x>
- Serbin, S. P., Singh, A., McNeil, B. E., Kingdon, C. C., & Townsend, P. A. (2014). Spectroscopic determination of leaf morphological and biochemical traits for northern temperate and boreal tree species. *Ecological Applications*, 24(7), 1651–1669. <https://doi.org/10.1890/13-2110.1>
- Ueda, Y., Siddique, S., & Frei, M. (2015). A novel gene, OZONE-RESPONSIVE APOPLASTIC PROTEIN1, enhances cell death in ozone stress in rice. *Plant Physiology*, 169(1), 873–889. <https://doi.org/10.1104/pp.15.00956>
- von Caemmerer, S., & Furbank, R. T. (2016). Strategies for improving C4 photosynthesis. *Current Opinion in Plant Biology*, 31, 125–134. <https://doi.org/10.1016/j.pbi.2016.04.003>
- Wang, J., Zeng, Q., Zhu, J., Chen, C., Liu, G., & Tang, H. (2014). Apoplastic antioxidant enzyme responses to chronic free-air ozone exposure in two different ozone-sensitive wheat cultivars. *Plant Physiology and Biochemistry*, 82, 183–193. <https://doi.org/10.1016/j.plaphy.2014.06.004>
- Wang, Y., Yang, S., Irfan, M., Zhang, C., Sun, Q., & Wu, S. (2013). Genetic analysis of carbon metabolism-related traits in maize using mixed major and polygene models. *Australian Journal of Crop Science*, 7(8), 1205–1211.
- Watanabe, M., Kamimaki, Y. U., Mori, M., Okabe, S., Arakawa, I., Kinose, Y., ... Izuta, T. (2018). Mesophyll conductance to CO_2 in leaves of Siebold's beech (*Fagus crenata*) seedlings under elevated ozone. *Journal of Plant Research*, 131(6), 907–914. <https://doi.org/10.1007/s10265-018-1063-4>
- Wayne, M. L., Telonis-Scott, M., Bono, L. M., Hasrchan, L., Kopp, A., Nuzhdin, S. V., & McIntyre, L. M. (2007). Simpler mode of inheritance of transcriptional variation in male *Drosophila melanogaster*. *Proceedings of the National Academy of Sciences of the United States of America*, 104(47), 18577–18582. <https://doi.org/10.1073/pnas.0705441104>
- Whitfield, A. C. P., Davison, A. W., & Ashenden, T. W. (1997). Artificial selection and heritability of ozone resistance in two populations of *Plantago major*. *New Phytologist*, 137(4), 645–655. <https://doi.org/10.1046/j.1469-8137.1997.00860.x>
- Wilkinson, S., & Davies, W. J. (2010). Drought, ozone, ABA and ethylene: New insights from cell to plant to community. *Plant Cell and Environment*, 33(4), 510–525. <https://doi.org/10.1111/j.1365-3040.2009.02052.x>
- Yendrek, C. R., Erice, G., Montes, C. M., Tomaz, T., Sorgini, C. A., Brown, P. J., ... Ainsworth, E. A. (2017). Elevated ozone reduces photosynthetic carbon gain by accelerating leaf senescence of

- inbred and hybrid maize in a genotype-specific manner. *Plant Cell and Environment*, 40(12), 3088–3100. <https://doi.org/10.1111/pce.13075>
- Yendrek, C. R., Tomaz, T., Montes, C. M., Cao, Y., Morse, A. M., Brown, P. J., ... Ainsworth, E. A. (2017). High-throughput phenotyping of maize leaf physiological and biochemical traits using hyperspectral reflectance. *Plant Physiology*, 173(1), 614–626. <https://doi.org/10.1104/pp.16.01447>
- Ziyomo, C., & Bernardo, R. (2013). Drought tolerance in maize: Indirect selection through secondary traits versus genomewide selection. *Crop Science*, 53(4), 1858–1870. <https://doi.org/10.2135/cropsci2012.11.0651>

SUPPORTING INFORMATION

Additional supporting information may be found online in the Supporting Information section at the end of the article.

How to cite this article: Choquette NE, Ogut F, Wertin TM, et al. Uncovering hidden genetic variation in photosynthesis of field-grown maize under ozone pollution. *Glob Change Biol.* 2019;00:1–12. <https://doi.org/10.1111/gcb.14794>